



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

hybrids, although attempts were persistently repeated throughout the summer. However, a few seeds were obtained from individuals having respectively *lata*, *gigas*, and intermediate number of chromosomes and the plants derived from these will form the chief subject of study for the coming year.

To summarize briefly:

The first generation offspring of *O. lata* ♀ × *O. gigas* ♂ fall into three main groups with respect to external characters and number of chromosomes; namely, *lata*, *gigas*-like and intermediate. Considering external characters only, the latter two should be further divided and subdivided.

Numbers of chromosomes are closely associated with external characters in the first and last, and probably also in the second group.

Pollen grains of two parental forms differ in number of lobes and these are inherited.

ANNE M. LUTZ

STATION FOR EXPERIMENTAL EVOLUTION,  
COLD SPRING HARBOR, L. I.,  
December 7, 1908

#### MUCOR CULTURES

In the study of the Mucoraceae for several years, some interesting facts concerning the development or rather the non-development of zygospores were observed. The experiments were made with the common *Mucor stolonifer* Ehrenberg. The media used were bread, pumpkin, orange, cornmeal, decoction of horse manure with gelatine, Pasteur's solution with gelatine, Hamaker culture medium.<sup>1</sup> The cultures were made with sterilized and unsterilized media. The spores for inoculation were taken from plants grown in the laboratory, from specimens collected for the herbarium, and from specimens sent to us by friends. In one thousand cultures not one zygospore was discovered.

In addition to the cultures, five hundred specimens of this species found growing spontaneously in different places were also examined but not one zygospore was observed.

Besides these experiments, many cultures were made and many specimens examined, a record of the exact number of which, however,

<sup>1</sup>Hamaker, SCIENCE, XXIII., 710, 1906.

was not kept. It is a conservative estimate to say that five hundred observations of this kind were made. This makes a grand total of two thousand observations without a single zygospore.

Experiments were also made to determine the development of this *Mucor* under anaerobic conditions. The media used for these experiments were orange, bread and Hamaker culture medium. All were sterilized. In giving the results of these experiments below, the word cornmeal will be used for the Hamaker medium. Cornmeal is the principal constituent of the medium. The material for inoculation was kindly furnished by Dr. Niewland, of Notre Dame University.

Small wide-mouthed bottles were used for the cultures. The medium was placed in the bottles and the bottles then closed with cotton and all sterilized. After inoculation, the bottles were placed into Novi jars and the jars filled with gas. The jars with the bottles were then set aside for observation. The following results were obtained.

*In Hydrogen.*—On orange, mycelium developed but few sporangiophores, no zygospores; on bread, no development; on cornmeal, no development.

*In Nitrogen.*—On orange, mycelium and few sporangiophores, no zygospores; on bread, about the same result; on cornmeal, about the same result.

*In Carbon Dioxide.*—On orange, mycelium well developed but few sporangiophores; on bread, mycelium profusely developed, many sporangiophores, no sporangia, no zygospores; on cornmeal, no development.

It seems that the absence of oxygen is not a necessary condition for the growth of zygospores.

DAVID R. SUMSTINE

UNIVERSITY OF PITTSBURG

#### THE BOTANICAL SOCIETY OF AMERICA

A UNION OF THE BOTANICAL SOCIETY OF AMERICA,  
THE SOCIETY FOR PLANT MORPHOLOGY AND  
PHYSIOLOGY AND THE AMERICAN  
MYCOLOGICAL SOCIETY

The third annual meeting of the federated societies (the fifteenth of the Botanical Society